

# Derivative-free optimization of composite functions

Jeffrey Larson

Argonne National Laboratory

May 27, 2019

# Problem setup

$$\begin{aligned} & \underset{x}{\text{minimize}} \quad f(x) = h(x; S(x)) \\ & \text{subject to: } x \in \mathcal{D} \subset \mathbb{R}^n \end{aligned}$$

where the objective  $f$  depends on the output(s) from a simulation  $S$  and a known function  $h$ .



# Problem setup


$$\begin{aligned} & \underset{x}{\text{minimize}} \quad f(x) = h(x; S(x)) \\ & \text{subject to: } x \in \mathcal{D} \subset \mathbb{R}^n \end{aligned}$$

where the objective  $f$  depends on the output(s) from a simulation  $S$  and a known function  $h$ .

- ▶ Derivatives of  $S$  may not be available
- ▶ Constraints defining  $\mathcal{D}$  may or may not depend on  $S$
- ▶ The dimension  $n$  is small
- ▶ Evaluating  $S$  is expensive (not using grids or randomized/evolutionary methods)



# Computers/Simulations!



# Computers/Simulations!




# Computers/Simulations!




#21 on TOP500 November 2018

(#6 on TOP500 June 2016)

# Model-based methods - Interpolation




# Model-based methods - Interpolation







# Model-based methods - Interpolation




# Model-based methods - Interpolation




# Model-based methods - Interpolation




# Model-based methods - Interpolation




# Model-based methods - Interpolation




# Model-based methods - Interpolation



# Model-based methods - Interpolation




# Model-based methods - Interpolation







# Model-based methods - Interpolation




## Model-based methods - Interpolation




## Model-based methods - Interpolation



## Model-based methods - Interpolation



## Model-based methods - Interpolation




# Opening up the black box

$$f(x) = \|S(x) - T\|_2^2 = \sum_{i=1}^p (S_i(x) - T_i)^2$$

Can either have a solver that uses  $f(x)$  or  $[S_1(x), \dots, S_p(x)]$ .



# Opening up the black box



Tuning quadrupole moments for a particle accelerator simulation.

$$f(x) = \|S(x) - T\|_2^2 = \sum_{i=1}^p (S_i(x) - T_i)^2$$

Can either have a solver that uses  $f(x)$  or  $[S_1(x), \dots, S_p(x)]$ .

# Emittance minimization

$$\underset{v \in \mathbb{R}^n}{\text{minimize}} \epsilon(v)$$

$$\text{subject to: } v \in \mathcal{D}$$

where

$$\epsilon(v) = \sqrt{\langle x(v)^2 \rangle \langle p_x(v)^2 \rangle - \langle x p_x(v) \rangle^2}$$






# Emittance minimization

$$\underset{v \in \mathbb{R}^n}{\text{minimize}} \epsilon(v)$$

$$\text{subject to: } v \in \mathcal{D}$$

where

$$\epsilon(v) = \sqrt{\langle x(v)^2 \rangle \langle p_x(v)^2 \rangle - \langle x p_x(v) \rangle^2}$$




# Emittance minimization

$$\begin{aligned} & \underset{v \in \mathbb{R}^n}{\text{minimize}} \quad \min_t \epsilon(v, t) \\ & \text{subject to: } v \in \mathcal{D} \\ & \quad \quad \quad 0 \leq t \leq \bar{t}, \end{aligned}$$

where

$$\epsilon(v, t) = \sqrt{\langle x(v, t)^2 \rangle \langle p_x(v, t)^2 \rangle - \langle x p_x(v, t) \rangle^2}$$



# Exploiting Structure

- ▶ Nonsmooth, composite optimization

$$\underset{x}{\text{minimize}} \ f(x) = h(S(x))$$


nonsmooth  $h: \mathbb{R}^p \rightarrow \mathbb{R}$  (with a known structure), smooth  $S: \mathbb{R}^n \rightarrow \mathbb{R}^p$  (expensive to evaluate).

- ▶ Idea: Build  $p$  models, one for each component of  $S$ . Use model gradients in place of  $\nabla S$ .
- ▶ Requires a *manifold representation* of  $h$ .
- ▶ Example: censored  $\ell_1$  loss:


$$f(x) = \sum_{i=1}^p |d_i - \max\{c_i, S_i(x)\}|$$



## Censored $\ell_1$ loss



## Censored $\ell_1$ loss



# Manifold representation

►  $h(y) = \max_{i \in \{1, \dots, p\}} y_i$

$p$  manifolds



# Manifold representation

►  $h(y) = \max_{i \in \{1, \dots, p\}} y_i$   $p$  manifolds

►  $h(y) = \|y\|_\infty = \max_{i \in \{1, \dots, p\}} |y_i|$   $2p$  manifolds



# Manifold representation

►  $h(y) = \max_{i \in \{1, \dots, p\}} y_i$   $p$  manifolds

►  $h(y) = \|y\|_\infty = \max_{i \in \{1, \dots, p\}} |y_i|$   $2p$  manifolds

►  $h(y) = \|y\|_1 = \sum_{i=1}^p |y_i|$   $2^p$  manifolds





# Manifold representation

►  $h(y) = \max_{i \in \{1, \dots, p\}} y_i$   $p$  manifolds

►  $h(y) = \|y\|_{\infty} = \max_{i \in \{1, \dots, p\}} |y_i|$   $2p$  manifolds

►  $h(y) = \|y\|_1 = \sum_{i=1}^p |y_i|$   $2^p$  manifolds

►  $h(y) = \sum_{i=1}^p |d_i - \max\{c_i, y_i\}|$   $3^p$  manifolds. If  $p = 45$ ,  
approximately  $3 \times 10^{21}$  potential manifolds.



# Manifold representation

►  $h(y) = \max_{i \in \{1, \dots, p\}} y_i$   $p$  manifolds

►  $h(y) = \|y\|_\infty = \max_{i \in \{1, \dots, p\}} |y_i|$   $2p$  manifolds

►  $h(y) = \|y\|_1 = \sum_{i=1}^p |y_i|$   $2^p$  manifolds


►  $h(y) = \sum_{i=1}^p |d_i - \max \{c_i, y_i\}|$   $3^p$  manifolds. If  $p = 45$ ,  
approximately  $3 \times 10^{21}$  potential manifolds.

User scripts need to calculate:


$f(x), S(x), \mathbb{H}(S(x)), \{\nabla h_i(S(x)) : i \in \mathbb{H}(S(x))\}, \{h_i(S(x)) : i \in \mathbb{G}\},$




# Manifold Sampling




# Manifold Sampling




# Manifold Sampling




# Manifold Sampling



# Manifold Sampling




# Manifold Sampling







# Manifold Sampling




# Manifold Sampling




# Manifold Sampling




# Manifold Sampling




# Manifold Sampling




# Manifold Sampling



# Manifold Sampling




# Manifold Sampling







# Manifold Sampling




# Manifold Sampling



# Manifold Sampling



# Manifold Sampling



# Smooth master model

$$g^k \triangleq \mathbf{proj} \left( 0, \mathbf{co} \left( \mathbb{G}^k \right) \right) \in \mathbf{co} \left( \mathbb{G}^k \right),$$



# Smooth master model

$$g^k \triangleq \mathbf{proj} (0, \mathbf{co} (\mathbb{G}^k)) \in \mathbf{co} (\mathbb{G}^k) ,$$

where

$$\mathbb{G}^k \triangleq \bigcup_{i \in I_h(S(x^k))} \{ \nabla M(x^k) \nabla h_i(S(x^k)) \}$$



# Smooth master model

$$g^k \triangleq \mathbf{proj} \left( 0, \mathbf{co} \left( \mathbb{G}^k \right) \right) \in \mathbf{co} \left( \mathbb{G}^k \right),$$

where

$$\mathbb{G}^k \triangleq \bigcup_{i \in I_h(S(x^k))} \{ \nabla M(x^k) \nabla h_i(S(x^k)) \}$$

or

$$\mathbb{G}^k \triangleq \bigcup_{y \in Y} \bigcup_{i \in I_h(S(y))} \{ \nabla M(x^k) \nabla h_i(S(x^k)) \}$$



# Smooth master model

$$g^k \triangleq \mathbf{proj} (0, \mathbf{co} (\mathbb{G}^k)) \in \mathbf{co} (\mathbb{G}^k) ,$$

where

$$\mathbb{G}^k \triangleq \bigcup_{i \in I_h(S(x^k))} \{ \nabla M(x^k) \nabla h_i(S(x^k)) \}$$

or

$$\mathbb{G}^k \triangleq \bigcup_{y \in Y} \bigcup_{i \in I_h(S(y))} \{ \nabla M(x^k) \nabla h_i(S(x^k)) \}$$


Define the smooth *master model*  $m_k^f: \mathbb{R}^n \rightarrow \mathbb{R}$  (with gradient  $g^k$ ) and obtain step by (approximately) solving

$$\begin{aligned} & \underset{s}{\text{minimize}} \quad m_k^f(x^k + s) \\ & \text{subject to: } s \in \mathcal{B}(0, \Delta_k) \end{aligned}$$







# Manifold Sampling




# Manifold Sampling




# Manifold Sampling




# Manifold Sampling




# Manifold Sampling




# Manifold Sampling



# Manifold Sampling




# Manifold Sampling







# Manifold Sampling




# Manifold Sampling




# Manifold Sampling




# Manifold Sampling




# Manifold Sampling




# Manifold Sampling



# Manifold Sampling




# Manifold Sampling







# Manifold Sampling




# Manifold Sampling




# Manifold Sampling




# Manifold Sampling




# Manifold Sampling




# Manifold Sampling



# Manifold Sampling



# Manifold Sampling





# Better trust-region subproblem?

Instead of solving

$$\begin{aligned} & \underset{s}{\text{minimize}} \quad m_k^f(x^k + s) \\ & \text{subject to: } s \in \mathcal{B}(0, \Delta_k) \end{aligned}$$

How about

$$\begin{aligned} & \underset{s}{\text{minimize}} \quad h(M(x^k + s)) \\ & \text{subject to: } s \in \mathcal{B}(0, \Delta_k) \end{aligned}$$



# Better trust-region subproblem?

Instead of solving

$$\begin{aligned} & \underset{s}{\text{minimize}} \quad m_k^f(x^k + s) \\ & \text{subject to: } s \in \mathcal{B}(0, \Delta_k) \end{aligned}$$

How about


$$\begin{aligned} & \underset{s}{\text{minimize}} \quad h(M(x^k + s)) \\ & \text{subject to: } s \in \mathcal{B}(0, \Delta_k) \end{aligned}$$

For censored  $\ell_1$  loss:


$$\begin{aligned} & \underset{s}{\text{minimize}} \quad \sum_{i=1}^p |d_i - \max \{c_i, q_i(x)\}| \\ & \text{subject to: } s \in \mathcal{B}(0, \Delta_k) \end{aligned}$$




# Manifold Sampling




# Manifold Sampling




# Manifold Sampling




# Manifold Sampling



# Manifold Sampling




# Manifold Sampling







# Manifold Sampling




# Manifold Sampling




# Manifold Sampling




# Manifold Sampling




# Manifold Sampling




# Manifold Sampling



# Manifold Sampling




# Manifold Sampling





# Manifold Sampling



# Better trust-region subproblem?

Instead of solving

$$\begin{aligned} & \underset{s}{\text{minimize}} \quad m_k^f(x^k + s) \\ & \text{subject to: } s \in \mathcal{B}(0, \Delta_k) \end{aligned}$$

How about

$$\begin{aligned} & \underset{s}{\text{minimize}} \quad h(M(x^k + s)) \\ & \text{subject to: } s \in \mathcal{B}(0, \Delta_k) \end{aligned}$$

For censored  $\ell_1$  loss:

$$\begin{aligned} & \underset{s}{\text{minimize}} \quad \sum_{i=1}^p |d_i - \max\{c_i, q_i(x)\}| \\ & \text{subject to: } s \in \mathcal{B}(0, \Delta_k) \end{aligned}$$



# Better trust-region subproblem?

Instead of solving

$$\begin{aligned} & \underset{s}{\text{minimize}} \quad m_k^f(x^k + s) \\ & \text{subject to: } s \in \mathcal{B}(0, \Delta_k) \end{aligned}$$

How about

$$\begin{aligned} & \underset{s}{\text{minimize}} \quad h(M(x^k + s)) \\ & \text{subject to: } s \in \mathcal{B}(0, \Delta_k) \end{aligned}$$

For censored  $\ell_1$  loss:

$$\begin{aligned} & \underset{s}{\text{minimize}} \quad \sum_{i=1}^p |d_i - \max \{c_i, q_i(x)\}| \\ & \text{subject to: } s \in \mathcal{B}(0, \Delta_k) \end{aligned}$$

## Question

Best method for solving composite nonsmooth quadratic problems?

# Thanks

Questions?

jmlarson@anl.gov

